Product Datasheet

CD20 Antibody (MS4A1/8072R) [Janelia Fluor® 525] NBP3-24187JF525

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-24187JF525

Updated 8/20/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-24187JF525



NBP3-24187JF525

Product Information	CD20 Antibody (MS4A1/8072R) [Janelia Fluor® 525]	
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone MS4A1/8072R Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Janelia Fluor 525 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues, Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Western Blot, Immunohistochemistry-Paraffin	Unit Size	0.1 ml
Clone MS4A1/8072R Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Janelia Fluor 525 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and Nymphoid tissues, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues, Rarely, CD2O-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions	Concentration	·
Clone MS4A1/8072R Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Janelia Fluor 525 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Rabbit Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in itssues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Notes Product Application Details Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Janelia Fluor 525 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid itssues, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody or ascribing a B-cell phenotype in known lymphoid tissues, Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Western Blot, Immunohistochemistry-Paraffin	Clonality	Monoclonal
IgG Kappa IgG Kappa	Clone	MS4A1/8072R
Conjugate Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and I will blood and	Preservative	0.05% Sodium Azide
Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and I ymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Western Blot, Immunohistochemistry-Paraffin	Isotype	IgG Kappa
Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Conjugate	Janelia Fluor 525
Product Description Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Western Blot, Immunohistochemistry-Paraffin	Purity	Protein A purified
Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Buffer	50mM Sodium Borate
Gene Symbol MS4A1 Species Human Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Product Description	
Gene Symbol MS4A1 Species Human B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Western Blot, Immunohistochemistry-Paraffin	Host	Rabbit
Species Human B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Gene ID	931
Marker B-Cell Marker Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Gene Symbol	MS4A1
Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Western Blot, Immunohistochemistry-Paraffin	Species	Human
malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa200-297) of human CD20 protein (exact sequence is proprietary) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Western Blot, Immunohistochemistry-Paraffin	Marker	B-Cell Marker
Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Specificity/Sensitivity	malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly
Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus. Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin Western Blot, Immunohistochemistry-Paraffin	Immunogen	
Applications Western Blot, Immunohistochemistry-Paraffin Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Notes	· ·
Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin	Product Application Details	
, , , , , , , , , , , , , , , , , , ,	Applications	Western Blot, Immunohistochemistry-Paraffin
Annlication Notes Ontimal dilution of this antibody should be experimentally determined	Recommended Dilutions	Western Blot, Immunohistochemistry-Paraffin
Optimal dilution of this antibody should be experimentally determined.	Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-24187JF525

NBP2-61455-10ug Recombinant Human CD20 His Protein

210-TA-005 TNF-alpha [Unconjugated]
NBL1-13312 CD20 Overexpression Lysate
AF114 CD45 Antibody [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-24187JF525

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

