## **Product Datasheet**

### Cytokeratin, pan Antibody (MonoPoly/4999R) [HRP] NBP3-24196H

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-24196H

Updated 10/7/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-24196H



#### NBP3-24196H

Cytokeratin, pan Antibody (MonoPoly/4999R) [HRP]

Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark.
Clonality	Monoclonal
Clone	MonoPoly/4999R
Preservative	No Preservative
Isotype	IgG Kappa
Conjugate	HRP
Purity	Protein A or G purified
Buffer	PBS
Product Description	
Host	Rabbit
Gene ID	3848
Gene Symbol	KRT1
Species	Human
Marker	Epithelial Marker
Specificity/Sensitivity	MonoPoly antibodies are designed by pooling several monospecific, recombinant monoclonal antibodies against a target. MonoPoly antibodies are a kind of synthetic polyclonal antibodies that can be produced in unlimited quantity with a strict lot-to-lot consistency. These antibodies are highly specific with exquisite sensitivity and a single MonoPoly antibody can be used for a variety of applications. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18), 40kDa (CK19) and 46kDa (CK20). This antibody is a broad-spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms.
Specificity/Sensitivity Immunogen	monoclonal antibodies against a target. MonoPoly antibodies are a kind of synthetic polyclonal antibodies that can be produced in unlimited quantity with a strict lot-to-lot consistency. These antibodies are highly specific with exquisite sensitivity and a single MonoPoly antibody can be used for a variety of applications. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18), 40kDa (CK19) and 46kDa (CK20). This antibody is a broad- spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during
	monoclonal antibodies against a target. MonoPoly antibodies are a kind of synthetic polyclonal antibodies that can be produced in unlimited quantity with a strict lot-to-lot consistency. These antibodies are highly specific with exquisite sensitivity and a single MonoPoly antibody can be used for a variety of applications. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18), 40kDa (CK19) and 46kDa (CK20). This antibody is a broad- spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. Recombinant fragments and/or synthetic peptides of human Cytokeratin, pan
Immunogen	monoclonal antibodies against a target. MonoPoly antibodies are a kind of synthetic polyclonal antibodies that can be produced in unlimited quantity with a strict lot-to-lot consistency. These antibodies are highly specific with exquisite sensitivity and a single MonoPoly antibody can be used for a variety of applications. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18), 40kDa (CK19) and 46kDa (CK20). This antibody is a broad- spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. Recombinant fragments and/or synthetic peptides of human Cytokeratin, pan
Immunogen Product Application Details	monoclonal antibodies against a target. MonoPoly antibodies are a kind of synthetic polyclonal antibodies that can be produced in unlimited quantity with a strict lot-to-lot consistency. These antibodies are highly specific with exquisite sensitivity and a single MonoPoly antibody can be used for a variety of applications. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18), 40kDa (CK19) and 46kDa (CK20). This antibody is a broad- spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. Recombinant fragments and/or synthetic peptides of human Cytokeratin, pan proteins (exact sequences are proprietary) Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,
Immunogen Product Application Details Applications	monoclonal antibodies against a target. MonoPoly antibodies are a kind of synthetic polyclonal antibodies that can be produced in unlimited quantity with a strict lot-to-lot consistency. These antibodies are highly specific with exquisite sensitivity and a single MonoPoly antibody can be used for a variety of applications. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 55kDa (CK7); 52kDa (CK8); 56.5kDa (CK10); 53kDa (CK13); 50kDa (CK14); 50kDa (CK15); 48kDa (CK16); 46kDa (CK17); 45kDa (CK18), 40kDa (CK19) and 46kDa (CK20). This antibody is a broad- spectrum anti pan-cytokeratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer. It is useful in characterizing the source of various neoplasms and to study the distribution of cytokeratin containing cells in epithelia during normal development and during the development of epithelial neoplasms. Recombinant fragments and/or synthetic peptides of human Cytokeratin, pan proteins (exact sequences are proprietary) Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-24196H

DNST0	Endostatin [HRP]
MAB1455	Albumin Antibody (188835) [Unconjugated] - Serum
NB100-687	Cytokeratin 19 Antibody - BSA Free
NBP2-16094	Cytokeratin 8 Antibody - BSA Free

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-24196H

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

