Product Datasheet

Cytokeratin 6 Antibody (rKRT6/9402) - Azide and BSA Free NBP3-24257

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-24257

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-24257



NBP3-24257

Cytokeratin 6 Antibody (rKRT6/9402) - Azide and BSA Free

Unit Size 100 ug Concentration 1 mg/ml Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone rKRT6/9402 Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Application Notes Immunohistochemistry-Paraffin I-2ug/ml Application Notes Immunohistochemistry-Paraffin 1-2ug/ml Immunohistochemistry-Paraffin I-2ug/ml Immunohistochemistry-Paraffin I-2ug/ml EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific applications hould be determined.	Cytokeratin 6 Antibody (rKR16/9402) - Azide and BSA Free	
Concentration 1 mg/ml Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone rKRT6/9402 Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires hearing tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Information	
Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone rKRT6/9402 Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Unit Size	100 ug
Clonality Monoclonal Clone rKRT6/9402 Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Concentration	1 mg/ml
Clone rKRT6/9402 Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Application Notes Immunohistochemistry-Paraffin I-2ug/ml Immunohistochemistry-Pa	Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml Immunohistochemistry-Paraffin: 1-2ug/ml Issue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype	Clone	rKRT6/9402
Purity Protein A or G purified Buffer 10mM PBS Product Description Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin 1-2ug/ml Recommended Dilutions Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	No Preservative
Buffer 10mM PBS Product Description Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG Kappa
Product Description Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin: 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
Description Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10mM PBS
Host Mouse Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID 3853 Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Antibody without azide - store at -20 to -80C.
Gene Symbol KRT6A Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
Species Human Marker Basal Cell Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	3853
Marker Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	KRT6A
Immunogen Recombinant fragment (around aa400-564) of human Cytokeratin 6 (exact sequence is proprietary) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Basal Cell Marker
Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	
Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry-Paraffin
formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry-Paraffin 1-2ug/ml
'	Application Notes	formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-24257

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NB100-355 RPE65 Antibody (401.8B11.3D9) - BSA Free

NB200-103 p53 Antibody (PAb 240) - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-24257

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

