Product Datasheet

Dicer Antibody (HL1232) - Azide and BSA Free NBP3-25439

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-25439

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-25439



NBP3-25439


Dicer Antibody (HL1232) - Azide and BSA Free

Dicer Antibody (HL1232) - Azide	and don fiee
Product Information	
Unit Size	100 ul
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	HL1232
Preservative	No Preservative
Isotype	IgG
Purity	Protein A purified
Buffer	PBS
Product Description	
Description	Novus Biologicals Rabbit Dicer Antibody (HL1232) - Azide and BSA Free (NBP3-25439) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	23405
Gene Symbol	DICER1
Species	Human, Mouse, Rat, Canine, Feline
Immunogen	Recombinant protein encompassing a sequence within the center region of human Dicer. The exact sequence is proprietary.
Product Application Details	
Applications	Western Plet Immunohistochemistry Paraffin Immunosytechemistry/

1 Todact Application Details	
	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
	Western Blot 1:500-1:3000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin


Images

Western Blot: Dicer Antibody (HL1232) - Azide and BSA Free [NBP3-25439] - Non-transfected (-) and transfected (+) MCF-7 whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with Dicer antibody [HL1232] (NBP3-25439) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.







Western Blot: Dicer Antibody (HL1232) - Azide and BSA Free [NBP3-25439] - Various whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with Dicer antibody [HL1232] (NBP3-25439) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Western Blot: Dicer Antibody (HL1232) - Azide and BSA Free [NBP3-25439] - Various whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with Dicer antibody [HL1232] (NBP3-25439) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: Dicer Antibody (HL1232) - Azide and BSA Free [NBP3-25439] - Dicer antibody [HL1232] detects Dicer protein at cytoplasm by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: Dicer stained by Dicer antibody [HL1232] (NBP3-25439) diluted at 1:500. Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (NBP2-43837) diluted at 1:1000. Blue: Fluoroshield with DAPI. Scale bar= 10um.




Western Blot: Dicer Antibody (HL1232) - Azide and BSA Free [NBP3-25439] - Various whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with Dicer antibody [HL1232] (NBP3-25439) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



Immunohistochemistry-Paraffin: Dicer Antibody (HL1232) - Azide and BSA Free [NBP3-25439] - Dicer antibody [HL1232] detects Dicer protein at cytoplasm by immunohistochemical analysis. Sample: Paraffinembedded mouse testis. Dicer stained by Dicer antibody [HL1232] (NBP3-25439) diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



Western Blot: Dicer Antibody (HL1232) - Azide and BSA Free [NBP3-25439] - Various whole cell extracts (30 ug) were separated by 5% SDS-PAGE, and the membrane was blotted with Dicer antibody [HL1232] (NBP3-25439) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-25439

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-88949PEP Dicer Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-25439

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

