

Product Datasheet

Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free NBP3-25558

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-25558

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-25558



NBP3-25558

Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free


Product Information	
Unit Size	100 ul
Concentration	Concentrations vary lot to lot. See vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	HL1471
Preservative	No Preservative
Isotype	IgG
Purity	Protein A purified
Buffer	PBS

Product Description	
Description	Novus Biologicals Rabbit Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free (NBP3-25558) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4157
Gene Symbol	MC1R
Species	Human, Mouse, Rat
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human Melanocortin-1 R/MC1R. The exact sequence is proprietary.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500-1:3000, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:1000

Images


Immunohistochemistry-Paraffin: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - MC1 Receptor antibody [HL1471] detects MC1 Receptor protein at cell membrane and cytoplasm by immunohistochemical analysis. Sample: Paraffin-embedded rat testis. MC1 Receptor stained by MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:400. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




Immunohistochemistry-Paraffin: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - MC1 Receptor antibody [HL1471] detects MC1 Receptor protein at cell membrane and cytoplasm by immunohistochemical analysis. Sample: Paraffin-embedded mouse testis. MC1 Receptor stained by MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:400. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




Western Blot: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - Non-transfected (-) and transfected (+) A375 whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunohistochemistry-Paraffin: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - MC1 Receptor antibody [HL1471] detects MC1 Receptor protein at cell membrane and cytoplasm by immunohistochemical analysis. Sample: Paraffin-embedded mouse kidney. MC1 Receptor stained by MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:400. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




Western Blot: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - Various extracts (50 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunohistochemistry-Paraffin: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - MC1 Receptor antibody [HL1471] detects MC1 Receptor protein at cell membrane and cytoplasm by immunohistochemical analysis. Sample: Paraffin-embedded rat kidney. MC1 Receptor stained by MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:400. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




Western Blot: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - Various tissue extracts (50 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Western Blot: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - Various whole cell extracts (30 ug) were separated by 12% SDS-PAGE, and the membrane was blotted with MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody was used to detect the primary antibody.




Immunocytochemistry/Immunofluorescence: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - MC1 Receptor antibody [HL1471] detects MC1 Receptor protein at cytoplasm by immunofluorescent analysis. Sample: A375 cells were fixed in ice-cold MeOH for 5 min. Green: MC1 Receptor stained by MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:500. Blue: Fluoroshield with DAPI .



Immunohistochemistry-Paraffin: Melanocortin-1 R/MC1R Antibody (HL1471) - Azide and BSA Free [NBP3-25558] - MC1 Receptor antibody [HL1471] detects MC1 Receptor protein at cell membrane by immunohistochemical analysis. Sample: Paraffin-embedded A375 xenograft . MC1 Receptor stained by MC1 Receptor antibody [HL1471] (NBP3-25558) diluted at 1:100. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



Frozen-sectioned mouse cerebellum. Green: MC1 Receptor stained by MC1 Receptor antibody [HL1471] diluted at 1:100. Blue: Fluoroshield with DAPI. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-25558

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00004157-Q01-10ug	Recombinant Human Melanocortin-1 R/MC1R GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-25558

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

