Product Datasheet

Cytokeratin 10 Antibody (18B6) NBP3-26178-100ul

Unit Size: 100 ul

Store at -20 to -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-26178

Updated 1/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-26178



NBP3-26178-100ul

Cytokeratin 10 Antibody (18B6)


Product Information	
Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20 to -70C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	18B6
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS, pH 7.4, 150mM NaCl, and 50% glycerol
Product Description	
Host	Rabbit
Gene ID	3858
Gene Symbol	KRT10
Species	Human
Immunogen	A synthesized peptide derived from Human Cytokeratin 10 [UniProt P13645]
Product Application Details	
Applications	ELISA, Immunohistochemistry
Recommended Dilutions	ELISA, Immunohistochemistry 1:50-1:200



Images

ntibody (1886) [NBP3-26178] -

Immunohistochemistry: Cytokeratin 10 Antibody (18B6) [NBP3-26178] -Image of Cytokeratin 10 Antibody (18B6) diluted at 1:50 and staining in paraffin-embedded human pancreati tissue performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.34% DAB.



Page 2 of 3 v.20.1 Updated 1/7/2025

Immunohistochemistry: Cytokeratin 10 Antibody (18B6) [NBP3-26178] -Image of Cytokeratin 10 Antibody (18B6) diluted at 1:50 and staining in paraffin-embedded human pancreati tissue performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.34% DAB.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-26178-100ul

H00003858-Q01-10ug	Recombinant Human Cytokeratin 10 GST (N-Term) Protein
NBP2-24891	Rabbit IgG Isotype Control
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-26178

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

