Product Datasheet

GDNF Antibody (11D2) NBP3-26556-100ul

Unit Size: 100 ul

Store at -20 to -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-26556

Updated 1/7/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-26556



NBP3-26556-100ul

GDNF Antibody (11D2)

| OBIN / Williams (1182) | |
|------------------------------------|---|
| Product Information | |
| Unit Size | 100 ul |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at -20 to -70C. Avoid freeze-thaw cycles. |
| Clonality | Monoclonal |
| Clone | 11D2 |
| Preservative | 0.02% Sodium Azide |
| Isotype | IgG |
| Purity | Affinity purified |
| Buffer | PBS, pH 7.4, 150mM NaCl, and 50% glycerol |
| Product Description | |
| Host | Rabbit |
| Gene ID | 2668 |
| Gene Symbol | GDNF |
| Species | Human |
| Immunogen | A synthesized peptide derived from Human GDNF [UniProt P39905] |
| Product Application Details | |
| Applications | Western Blot, ELISA, Flow Cytometry, Immunohistochemistry, Immunoprecipitation |
| Recommended Dilutions | Western Blot 1:500-1:5000, Flow Cytometry, ELISA, Immunohistochemistry 1:50 |

-1:200, Immunoprecipitation 1:200-1:1000

Images


Immunoprecipitation: GDNF Antibody (11D2) [NBP3-26556] -

Immunoprecipitating GDNF in Hela whole cell lysate.


Lane 1: Rabbit control IgG instead of NBP3-26556 in Hela whole cell lysate. For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000).

Lane 2: NBP3-26556 (3ug) + Hela whole cell lysate (500ug).


Lane 3: Hela whole cell lysate (20ug).



Immunohistochemistry: GDNF Antibody (11D2) [NBP3-26556] - Image of GDNF Antibody (11D2) diluted at 1:107.5 and staining in paraffinembedded human kidney tissue performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunohistochemistry: GDNF Antibody (11D2) [NBP3-26556] - Image of GDNF Antibody (11D2) diluted at 1:107.5 and staining in paraffinembedded human kidney tissue performed. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.




Western Blot: GDNF Antibody (11D2) [NBP3-26556] - Positive Western Blot detected in: HepG2 whole cell lysate, 293T whole cell lysate, SH-SY5Y whole cell lysate, PC3 whole cell lysate, Hela whole cell lysate, MCF-7 whole cell lysate, U251 whole cell lysate.

All lanes: GDNF Antibody at 1ug/ml.


Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution.

Predicted band size: 24, 21, 26, 23, 19 KDa

Observed band size: 24 KDa



Flow Cytometry: GDNF Antibody (11D2) [NBP3-26556] - Overlay histogram showing SH-SY5Y cells stained with GDNF Antibody (11D2) (red line) at 1:50. The cells were fixed with 70% Ethylalcohol (18h) and then permeabilized with 0.3% Triton X-100 for 2 min. The cells were then incubated in 1x PBS /10% normal goat serum to block non-specific protein-protein interactions followed by primary antibody for 1 h at 4C. The secondary antibody used was FITC goat anti-rabbit IgG (H+L) at 1/200 dilution for 1 h at 4C. Control antibody (green line) was used under the same conditions. Acquisition of >10,000 events was performed.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-26556-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-34888-10ug Recombinant Human GDNF Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-26556

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

