Product Datasheet

Biotin Antibody (rBTN/8819) - Azide and BSA Free NBP3-27043

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-27043

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-27043



Biotin Antibody (rBTN/8819) - Azi	de and BSA Free	
Product Information		
Unit Size	100 ug	
Concentration	1 mg/ml	
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.	
Clonality	Monoclonal	
Clone	rBTN/8819	
Preservative	No Preservative	
Isotype	IgG1 Kappa	
Purity	Protein A or G purified	
Buffer	10mM PBS	
Product Description		
Description	Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required.	
Host	Mouse	
Species	Non-species specific	
Specificity/Sensitivity	This antibody recognizes both the free and protein-conjugated (either soluble or cell bound) form of biotin. It is highly specific to biotin and shows no cross-reaction with other structurally related compounds. It has a very high affinity for biotin and is excellent for use in various amplification techniques. In some applications, localization of biotinylated probes with avidin produces unacceptably high background staining. Anti-biotin antibody may be substituted to decrease this noise.	
Immunogen	Biotinylated sheep immunoglobulin	
Product Application Details		
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, In-situ Hybridization	
Recommended Dilutions	Western Blot 0.5-2ug/ml, Flow Cytometry 0.5-1ug/million cells, Immunocytochemistry/ Immunofluorescence 0.5-1ug/ml, Immunohistochemistry-Paraffin 0.5-1ug/ml for 30 minutes at RT, In-situ Hybridization	
Application Notes	Immunohistochemistry-Paraffin 0.5-1ug/ml for 30 minutes at RT. Optimal dilution for a specific application should be determined Staining of formalin-fixed tissues	



min at 95C followed by cooling at RT for 20 minutes

requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45

Images	
Formalin-fixed, paraffin-embedded human kidney stained with Biotinylated Lambda Light Chain probe followed by Biotin Antibody (rBTN/8819) - Azide and BSA Free.	×





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-27043

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-27043

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

