

Product Datasheet

5'-Nucleotidase/CD73 Antibody (A6G2-R) NBP3-32166

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32166

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32166



NBP3-32166**5'-Nucleotidase/CD73 Antibody (A6G2-R)**

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	A6G2-R
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	63 kDa

Product Description	
Description	Novus Biologicals Mouse 5'-Nucleotidase/CD73 Antibody (A6G2-R) (NBP3-32166) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	4907
Gene Symbol	NT5E
Species	Human, Mouse, Rat
Immunogen	Recombinant protein within Human 5'-Nucleotidase/CD73 27-549. (Uniprot: P21589)

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1:100, Immunohistochemistry-Paraffin 1:1000



Images

Western Blot: 5'-Nucleotidase/CD73 Antibody (A6G2-R) [NBP3-32166] - Western blot analysis of 5'-Nucleotidase/CD73 on different lysates with Mouse anti-5'-Nucleotidase/CD73 antibody (NBP3-32166) at 1/2,000 dilution.

Lane 1: U-87 MG cell lysate

Lane 2: U-87 MG cell lysate (no heat)

Lysates/proteins at 30 ug/Lane.

Predicted band size: 63 kDa

Observed band size: 70 kDa

Exposure time: 24 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32166) at 1/2,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Anti-Mouse IgG for IP Nano-secondary antibody (NBI02H) at 1/5,000 dilution was used for 1 hour at room temperature.




Immunohistochemistry: 5'-Nucleotidase/CD73 Antibody (A6G2-R) [NBP3-32166] - Immunohistochemical analysis of paraffin-embedded NCI-H441 xenograft tissue with Mouse anti-5'-Nucleotidase/CD73 antibody (NBP3-32166) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (NBP3-32166) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Immunocytochemistry/ Immunofluorescence: 5'-Nucleotidase/CD73 Antibody (A6G2-R) [NBP3-32166] - Immunocytochemistry analysis of U-87 MG cells labeling 5'-Nucleotidase/CD73 with Mouse anti-5'-Nucleotidase/CD73 antibody (NBP3-32166) at 1/100 dilution.

Cells were fixed in 100% precooled methanol for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-5'-Nucleotidase/CD73 antibody (NBP3-32166) at 1/100 dilution in 1% BSA in PBST overnight at 4°C. Goat Anti-Mouse IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 594) were used as the secondary antibody at 1/1,000 dilution.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-32166

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBP2-51548-0.05mg	Recombinant Human 5'-Nucleotidase/CD73 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32166

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

