Product Datasheet

c-Myc Antibody (PS00-11) NBP3-32201

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32201

Updated 9/11/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-32201



NBP3-32201

c-Myc Antibody (PS00-11)

, , , , , , , , , , , , , , , , , , ,	
Product Information	
100 ul	
1 mg/ml	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Monoclonal	
PS00-11	
0.05% Sodium Azide	
IgG	
Protein A purified	
PBS (pH7.4), 0.1% BSA and 40% Glycerol	
49 kDa	
Product Description	
Rabbit	
4609	
MYC	
Human	
Synthetic peptide. (Uniprot: P01106)	
Product Application Details	
Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin	
Western Blot 1:1000, Immunohistochemistry, Immunohistochemistry-Paraffin 1:200	



Images

Western Blot: c-Myc Antibody (PS00-11) [NBP3-32201] - Western blot analysis of c-Myc on different lysates with Rabbit anti-c-Myc antibody (NBP3-32201) at 1/1,000 dilution.

Lane 1: Jurkat cell lysate Lane 2: Daudi cell lysate Lane 3: K562 cell lysate Lane 4: HL-60 cell lysate

Lysates/proteins at 10 ug/Lane.

Predicted band size: 49 kDa Observed band size: 55 kDa


Exposure time: 2 minutes;


10% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32201) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1:300,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: c-Myc Antibody (PS00-11) [NBP3-32201] - Immunohistochemical analysis of paraffin-embedded human B lymphocytoma tissue with Mouse anti-c-Myc antibody (NBP3-32201) at 1/200 dilution.

The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32201) at 1/200 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.









Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32201

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00004609-P01-10ug Recombinant Human c-Myc GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32201

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

