Product Datasheet

Cyclophilin 40 Antibody (PSH02-71) NBP3-32238

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32238

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-32238



NBP3-32238

Cyclophilin 40 Antibody (PSH02-71)

Cyclophilin 40 Antibody (PSH02-71)	
Product Information	
100 ul	
1 mg/ml	
Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.	
Monoclonal	
PSH02-71	
0.05% Sodium Azide	
IgG	
Protein A purified	
1*TBS (pH7.4), 0.1% BSA and 40% Glycerol	
47 kDa	
Product Description	
Novus Biologicals Rabbit Cyclophilin 40 Antibody (PSH02-71) (NBP3-32238) is a recombinant monoclonal antibody validated for use in IHC, WB, Flow and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.	
Rabbit	
5481	
PPID	
Human, Monkey	
Recombinant protein within human Cyclophilin 40 aa 151-370 / 370. (Uniprot: Q08752)	
Product Application Details	
Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry	
Western Blot 1:1000, Flow Cytometry 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100, Immunohistochemistry-Paraffin 1:2000	



Images

Western Blot: Cyclophilin 40 Antibody (PSH02-71) [NBP3-32238] - Western blot analysis of Cyclophilin 40 on different lysates with Rabbit anti-Cyclophilin 40 antibody (NBP3-32238) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: 293T cell lysate Lane 3: K-562 cell lysate

Lane 4: HepG2 cell lysate

Lane 5: MCF7 cell lysate

Lane 6: Jurkat cell lysate

Lane 7: COS-1 cell lysate

Lane 8: Human liver tissue lysate

Lysates/proteins at 30 ug/Lane.

Predicted band size: 41 kDa Observed band size: 47 kDa


Exposure time: 1 minutes 2 seconds;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32238) at 1/1,000 dilution was used in 5% NFDM/TBST at 4C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: Cyclophilin 40 Antibody (PSH02-71) [NBP3-32238] - Immunohistochemical analysis of paraffin-embedded human kidney tissue with Rabbit anti-Cyclophilin 40 antibody (NBP3-32238) at 1/2,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32238) at 1/2,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.








Immunocytochemistry/ Immunofluorescence: Cyclophilin 40 Antibody (PSH02-71) [NBP3-32238] - Immunocytochemistry analysis of 293T cells labeling Cyclophilin 40 with Rabbit anti-Cyclophilin 40 antibody (NBP3-32238) at 1/100 dilution.


Cells were fixed in 100% precooled methanol for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-Cyclophilin 40 antibody (NBP3-32238) at 1/100 dilution in 1% BSA in PBST overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4□. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.

Flow Cytometry: Cyclophilin 40 Antibody (PSH02-71) [NBP3-32238] - Flow cytometric analysis of 293T cells labeling Cyclophilin 40.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32238, 1:100) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4□ for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4□. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).









Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32238

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBC1-18427 Recombinant Human Cyclophilin 40 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32238

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

