Product Datasheet

alpha Glucosidase 2 Antibody (PSH01-53) NBP3-32367

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32367

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-32367



NBP3-32367

alpha Glucosidase 2 Antibody (PSH01-53)

alpha Glucosidase 2 Antibody (PSH01-53)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH01-53
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	107 kDa
Product Description	
Description	Novus Biologicals Rabbit alpha Glucosidase 2 Antibody (PSH01-53) (NBP3-32367) is a recombinant monoclonal antibody validated for use in IHC and WB. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	23193
Gene Symbol	GANAB
Species	Human
Immunogen	Recombinant protein within human alpha Glucosidase 2 aa 29-450. (Uniprot: Q14697)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
	Western Blot, mindrensteenermenty Faranin, mindrensteenermenty

	·
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry, Immunohistochemistry-Paraffin 1:1000



Images

Western Blot: alpha Glucosidase 2 Antibody (PSH01-53) [NBP3-32367] - Western blot analysis of alpha Glucosidase 2 on different lysates with Rabbit anti-alpha Glucosidase 2 antibody (NBP3-32367) at 1/1,000 dilution.

Lane 1: HeLa cell lysate (20 ug/Lane)

Lane 2: HepG2 cell lysate (20 ug/Lane)

Lane 3: LoVo cell lysate (20 ug/Lane)

Lane 4: HEK-293 cell lysate (20 ug/Lane)

Lane 5: U-2 OS cell lysate (20 ug/Lane)

Lane 6: K-562 cell lysate (20 ug/Lane)

Lane 7: Jurkat cell lysate (20 ug/Lane)

Lane 8: HL-60 cell lysate (20 ug/Lane)

Lane 9: Raji cell lysate (20 ug/Lane)

Predicted band size: 107 kDa Observed band size: 107 kDa

Exposure time: 1 minute 2 seconds;

4-20% SDS-PAGE gel.


Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32367) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.


Immunohistochemistry: alpha Glucosidase 2 Antibody (PSH01-53) [NBP3-32367] - Immunofluorescence analysis of paraffin-embedded human thyroid tissue labeling alpha Glucosidase 2 with Rabbit anti-alpha Glucosidase 2 antibody (NBP3-32367) at 1/200 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 10% negative goat serum for 1 hour at room temperature, washed with PBS, and then probed with the primary antibody (NBP3-32367, green) at 1/200 dilution overnight at 4 □, washed with PBS. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. Nuclei were counterstained with DAPI (blue).

Immunohistochemistry: alpha Glucosidase 2 Antibody (PSH01-53) [NBP3-32367] - Immunohistochemical analysis of paraffin-embedded human liver tissue with Rabbit anti-alpha Glucosidase 2 antibody (NBP3-32367) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32367) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.











Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32367

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00023193-P01-10ug Recombinant Human alpha Glucosidase 2 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32367

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

