

Product Datasheet

GRP75/HSPA9B/Mortalin Antibody (6-A9-R) NBP3-32407

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32407

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32407



NBP3-32407

GRP75/HSPA9B/Mortalin Antibody (6-A9-R)

Product Information

Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	6-A9-R
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	74 kDa

Product Description

Description	Novus Biologicals Mouse GRP75/HSPA9B/Mortalin Antibody (6-A9-R) (NBP3-32407) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Mouse
Gene ID	3313
Gene Symbol	HSPA9
Species	Human, Mouse, Rat
Immunogen	Synthetic peptide within human GRP75/HSPA9B/Mortalin aa 101-150. (Uniprot: P38646)

Product Application Details

Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:2000, Immunohistochemistry, Immunocytochemistry/Immunofluorescence 1:100, Immunohistochemistry-Paraffin 1:500



Images

Western Blot: GRP75/HSPA9B/Mortalin Antibody (6-A9-R) [NBP3-32407] - Western blot analysis of GRP75/HSPA9B/Mortalin on different lysates with Mouse anti-GRP75/HSPA9B/Mortalin antibody (NBP3-32407) at 1/2,000 dilution.

Lane 1: HeLa cell lysate
 Lane 2: K-562 cell lysate
 Lane 3: HepG2 cell lysate
 Lane 4: MCF7 cell lysate
 Lane 5: NIH/3T3 cell lysate
 Lane 6: Mouse liver tissue lysate
 Lane 7: Rat liver tissue lysate
 Lane 8: Mouse kidney tissue lysate
 Lane 9: Rat kidney tissue lysate

Predicted band size: 74 kDa
 Observed band size: 74 kDa



Exposure time: 3 minutes;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32407) at 1/2,000 dilution was used in 5% NFDM/TBST at 4 overnight. Anti-Mouse IgG for IP Nano-secondary antibody (NBI02H) at 1/5,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: GRP75/HSPA9B/Mortalin Antibody (6-A9-R) [NBP3-32407] - Immunohistochemical analysis of paraffin-embedded human liver tissue with Mouse anti-GRP75/HSPA9B/Mortalin antibody (NBP3-32407) at 1/500 dilution.


The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0) for 2 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (NBP3-32407) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.



Immunocytochemistry/ Immunofluorescence: GRP75/HSPA9B/Mortalin Antibody (6-A9-R) [NBP3-32407] - Immunocytochemistry analysis of K-562 cells labeling GRP75/HSPA9B/Mortalin with Mouse anti-GRP75/HSPA9B/Mortalin antibody (NBP3-32407) at 1/100 dilution.

Cells were fixed in 80% precooled methanol for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Mouse anti-GRP75/HSPA9B/Mortalin antibody (NBP3-32407) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

beta Tubulin (red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 594) were used as the secondary antibody at 1/1,000 dilution.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-32407

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBC1-18380	Recombinant Human GRP75/HSPA9B/Mortalin His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32407

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

