

# Product Datasheet

## LIAS Antibody (PSH04-04)

### NBP3-32536

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP3-32536](http://www.novusbio.com/NBP3-32536)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP3-32536](http://www.novusbio.com/reviews/destination/NBP3-32536)



**NBP3-32536**

LIAS Antibody (PSH04-04)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH04-04
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	42 kDa
Product Description	
Description	Novus Biologicals Rabbit LIAS Antibody (PSH04-04) (NBP3-32536) is a recombinant monoclonal antibody validated for use in WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	11019
Gene Symbol	LIAS
Species	Human, Mouse, Rat
Immunogen	Recombinant protein within human LIAS aa 1-372 / 372. (Uniprot: O43766)
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:2000, Immunocytochemistry/ Immunofluorescence 1:100



## Images

Western Blot: LIAS Antibody (PSH04-04) [NBP3-32536] - Western blot analysis of LIAS on different lysates with Rabbit anti-LIAS antibody (NBP3-32536) at 1/2,000 dilution.

Lane 1: K-562 cell lysate (20 ug/Lane)  
 Lane 2: LO2 cell lysate (20 ug/Lane)  
 Lane 3: Raji cell lysate (20 ug/Lane)  
 Lane 4: MCF7 cell lysate (20 ug/Lane)  
 Lane 5: Jurkat cell lysate (20 ug/Lane)  
 Lane 6: HepG2 cell lysate (20 ug/Lane)  
 Lane 7: PC-3M cell lysate (20 ug/Lane)  
 Lane 8: C2C12 cell lysate (20 ug/Lane)  
 Lane 9: L6 cell lysate (20 ug/Lane)  
 Lane 10: Mouse liver tissue lysate (40 ug/Lane)  
 Lane 11: Mouse colon tissue lysate (40 ug/Lane)  
 Lane 12: Mouse brain tissue lysate (40 ug/Lane)  
 Lane 13: Rat liver tissue lysate (40 ug/Lane)  
 Lane 14: Rat testis tissue lysate (40 ug/Lane)

Predicted band size: 42 kDa  
 Observed band size: 39 kDa



Exposure time: 42 seconds; ECL;  
 4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32536) at 1/2,000 dilution was used in 5% NFDM/TBST at 4 overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.

Immunocytochemistry/ Immunofluorescence: LIAS Antibody (PSH04-04) [NBP3-32536] - Immunocytochemistry analysis of C2C12 cells labeling LIAS with Rabbit anti-LIAS antibody (NBP3-32536) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-LIAS antibody (NBP3-32536) at 1/100 dilution in 1% BSA in PBST overnight at 4 °C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4 °C. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP3-32536**

---

NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP3-32536](http://www.novusbio.com/reviews/submit/NBP3-32536)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

