Product Datasheet

NDUFS2 Antibody (PSH03-26) NBP3-32639

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-32639

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-32639



NBP3-32639

NDUFS2 Antibody (PSH03-26)

NDUFS2 Antibody (PSHU3-26)	
Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH03-26
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	53 kDa
Product Description	
Description	Novus Biologicals Rabbit NDUFS2 Antibody (PSH03-26) (NBP3-32639) is a recombinant monoclonal antibody validated for use in IHC, WB, Flow, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4720
Gene Symbol	NDUFS2
Species	Human, Mouse, Rat
Immunogen	Recombinant protein within human NDUFS2 aa 1-463 / 463. (Uniprot: O75306)
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:2000, Flow Cytometry 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100, Immunoprecipitation 1-

2ug/sample, Immunohistochemistry-Paraffin 1:1000

Images

Immunoprecipitation: NDUFS2 Antibody (PSH03-26) [NBP3-32639] - NDUFS2 was immunoprecipitated in 0.2mg HeLa cell lysate with NBP3-32639 at 2 ug/25 ul agarose. Western blot was performed from the immunoprecipitate using NBP3-32639 at 1/2,000 dilution. Anti-Rabbit IgG for IP Nano-secondary antibody at 1/5,000 dilution was used for 1 hour at room temperature.




Lane 2: NBP3-32639 IP in HeLa cell lysate

Lane 3: Rabbit IgG instead of NBP3-32639 in HeLa cell lysate

Blocking/Dilution buffer: 5% NFDM/TBST

Exposure time: 43 seconds





Western Blot: NDUFS2 Antibody (PSH03-26) [NBP3-32639] - Western blot analysis of NDUFS2 on different lysates with Rabbit anti-NDUFS2 antibody (NBP3-32639) at 1/2,000 dilution.

Lane 1: HeLa cell lysate (20 ug/Lane)

Lane 2: Jurkat cell lysate (20 ug/Lane)

Lane 3: 293T cell lysate (20 ug/Lane)

Lane 4: A431 cell lysate (20 ug/Lane)

Lane 5: Raji cell lysate (20 ug/Lane)

Lane 6: SK-Br-3 cell lysate (20 ug/Lane)

Lane 7: K-562 cell lysate (20 ug/Lane)

Lane 8: NIH/3T3 cell lysate (20 ug/Lane)

Lane 9: RAW264.7 cell lysate (20 ug/Lane)

Lane 10: PC-12 cell lysate (20 ug/Lane)

Lane 11: Mouse brain tissue lysate (40 ug/Lane)

Lane 12: Mouse kidney tissue lysate (40 ug/Lane)

Lane 13: Rat brain tissue lysate (40 ug/Lane)

Lane 14: Rat kidney tissue lysate (40 ug/Lane)

Predicted band size: 53 kDa Observed band size: 45 kDa


Exposure time: 24 seconds;


4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32639) at 1/2,000 dilution was used in 5% NFDM/TBST at 4 overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.

Immunohistochemistry: NDUFS2 Antibody (PSH03-26) [NBP3-32639] - Immunohistochemical analysis of paraffin-embedded human kidney tissue with Rabbit anti-NDUFS2 antibody (NBP3-32639) at 1/1,000 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-32639) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.








Immunocytochemistry/ Immunofluorescence: NDUFS2 Antibody (PSH03 -26) [NBP3-32639] - Immunocytochemistry analysis of A431 cells labeling NDUFS2 with Rabbit anti-NDUFS2 antibody (NBP3-32639) at 1/100 dilution.


Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-NDUFS2 antibody (NBP3-32639) at 1/100 dilution in 1% BSA in PBST overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4□. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.

Flow Cytometry: NDUFS2 Antibody (PSH03-26) [NBP3-32639] - Flow cytometric analysis of A431 cells labeling NDUFS2.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32639, 1µg/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4□ for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4□. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).









Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-32639

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-23265 Recombinant Human NDUFS2 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32639

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

