

Product Datasheet

IRE1 [p Ser724] Antibody (PSH03-35) NBP3-32774

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-32774

Updated 8/7/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-32774



NBP3-32774

IRE1 [p Ser724] Antibody (PSH03-35)

Product Information	
Unit Size	100 ul
Concentration	1 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	PSH03-35
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Protein A purified
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol
Target Molecular Weight	110 kDa

Product Description	
Host	Rabbit
Gene ID	2081
Gene Symbol	ERN1
Species	Human, Mouse
Immunogen	Synthetic phospho-peptide corresponding to residues surrounding Ser724 of human IRE1 alpha. (Uniprot: O75460)

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:500, Immunohistochemistry-Paraffin 1:500



Images

Western Blot: IRE1 [p Ser724] Antibody (PSH03-35) [NBP3-32774] - Western blot analysis of IRE1 on different lysates with Rabbit anti-IRE1 antibody (NBP3-32774) at 1/1,000 dilution.

Lane 1: HeLa cell lysate

Lane 2: HeLa starved for 3 hours then treated with 100nM Calyculin A for 30 minutes cell lysate

Lane 3: Jurkat cell lysate

Lane 4: Jurkat treated with 100nM Calyculin A for 30 minutes cell lysate

Lysates/proteins at 20 ug/Lane.

Predicted band size: 54 kDa

Observed band size: 54 kDa

Exposure time: 5 minutes 10 seconds;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-32774) at 1/1,000 dilution was used in 5% NFDM/TBST at 4°C overnight. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.




Immunohistochemistry: IRE1 [p Ser724] Antibody (PSH03-35) [NBP3-32774] - Immunohistochemical analysis of paraffin-embedded human pancreas tissue with Rabbit anti-IRE1 antibody (NBP3-32774) at 1/500 dilution.

The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH₂O and PBS, and then probed with the primary antibody (NBP3-32774) at 1/500 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.

Immunocytochemistry/ Immunofluorescence: IRE1 [p Ser724] Antibody (PSH03-35) [NBP3-32774] - Immunocytochemistry analysis of Jurkat cells treated with 100nM Calyculin A for 30 minutes labeling IRE1 with Rabbit anti-IRE1 antibody (NBP3-32774) at 1/500 dilution.


Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-IRE1 antibody (NBP3-32774) at 1/500 dilution in 1% BSA in PBST overnight at 4°C. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4°C. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.



Flow Cytometry: IRE1 [p Ser724] Antibody (PSH03-35) [NBP3-32774] - Flow cytometric analysis of HeLa cells treated with or without 100nM Calyculin A for 30 minutes labeling IRE1.

Cells were fixed and permeabilized. Then stained with the primary antibody (NBP3-32774, 1µg/mL) (red) compared with Rabbit IgG Isotype Control (green). After incubation of the primary antibody at +4h for an hour, the cells were stained with a iFluor™ 488 conjugate-Goat anti-Rabbit IgG Secondary antibody at 1/1,000 dilution for 30 minutes at +4h. Unlabelled sample was used as a control (cells without incubation with primary antibody; black).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-32774

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

