Product Datasheet

AP2 alpha Antibody (PSH02-06) NBP3-33044

Unit Size: 100 ul

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-33044

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-33044



NBP3-33044

AP2 alpha Antibody (PSH02-06)

AFZ alpha Ahilibudy (F31102-00	o)			
Product Information				
Unit Size	100 ul			
Concentration	1 mg/ml			
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.			
Clonality	Monoclonal			
Clone	PSH02-06			
Preservative	0.05% Sodium Azide			
Isotype	IgG			
Purity	Protein A purified			
Buffer	PBS (pH7.4), 0.1% BSA and 40% Glycerol			
Target Molecular Weight	48 kDa			
Product Description				
Description	Novus Biologicals Rabbit AP2 alpha Antibody (PSH02-06) (NBP3-33044) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.			
Host	Rabbit			
Gene ID	7020			
Gene Symbol	TFAP2A			
Species	Human			
Immunogen	Recombinant protein within human AP2 alpha aa 1-437 / 437. (Uniprot: P05549)			
Product Application Details				
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry			
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 1:100, Immunohistochemistry-Paraffin 1:1000			

Images

Western Blot: AP2 alpha Antibody (PSH02-06) [NBP3-33044] - Western blot analysis of AP2 alpha on different lysates with Rabbit anti-AP2 alpha antibody (NBP3-33044) at 1/1,000 dilution.

Lane 1: HepG2 cell lysate Lane 2: SiHa cell lysate

Lysates/proteins at 20 ug/Lane.

Predicted band size: 48 kDa Observed band size: 48 kDa

Exposure time: 5 minutes;

4-20% SDS-PAGE gel.

Proteins were transferred to a PVDF membrane and blocked with 5% NFDM/TBST for 1 hour at room temperature. The primary antibody (NBP3-33044) at 1/1,000 dilution was used in 5% NFDM/TBST at room temperature for 2 hours. Goat Anti-Rabbit IgG - HRP Secondary Antibody at 1/50,000 dilution was used for 1 hour at room temperature.


Immunohistochemistry: AP2 alpha Antibody (PSH02-06) [NBP3-33044] - Immunohistochemical analysis of paraffin-embedded human breast tissue with Rabbit anti-AP2 alpha antibody (NBP3-33044) at 1/1,000 dilution.


The section was pre-treated using heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 20 minutes. The tissues were blocked in 1% BSA for 20 minutes at room temperature, washed with ddH2O and PBS, and then probed with the primary antibody (NBP3-33044) at 1/1,000 dilution for 1 hour at room temperature. The detection was performed using an HRP conjugated compact polymer system. DAB was used as the chromogen. Tissues were counterstained with hematoxylin and mounted with DPX.


Immunocytochemistry/ Immunofluorescence: AP2 alpha Antibody (PSH02-06) [NBP3-33044] - Immunocytochemistry analysis of HepG2 cells labeling AP2 alpha with Rabbit anti-AP2 alpha antibody (NBP3-33044) at 1/100 dilution.

Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 5 minutes at room temperature, then blocked with 1% BSA in 10% negative goat serum for 1 hour at room temperature. Cells were then incubated with Rabbit anti-AP2 alpha antibody (NBP3-33044) at 1/100 dilution in 1% BSA in PBST overnight at 4 □. Goat Anti-Rabbit IgG H&L (iFluor™ 488) was used as the secondary antibody at 1/1,000 dilution. PBS instead of the primary antibody was used as the secondary antibody only control. Nuclear DNA was labelled in blue with DAPI.

Beta tubulin (red) was stained at 1/100 dilution overnight at +4□. Goat Anti-Mouse IgG H&L (iFluor™ 594) was used as the secondary antibody at 1/1,000 dilution.











Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-33044

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00007020-P01-10ug Recombinant Human AP2 alpha GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-33044

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

