

Product Datasheet

Cysteinyl Leukotriene R1/CysLTR1 Antibody - BSA Free NLS1317

Unit Size: 0.05 ml

Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NLS1317

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NLS1317



NLS1317**Cysteinyl Leukotriene R1/CysLTR1 Antibody - BSA Free**

Product Information	
Unit Size	0.05 ml
Concentration	1.0 mg/ml
Storage	Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS


Product Description	
Description	Product can be stored undiluted at 4C for up to 1 month.
Host	Rabbit
Gene ID	10800
Gene Symbol	CYSLTR1
Species	Human, Monkey, Primate
Reactivity Notes	Predicted cross-reactivity based on sequence identity: Gorilla (100%), Gibbon (100%), Rabbit (85%), Guinea (85%), Bovine (80%), Equine (80%), Porcine (80%).
Specificity/Sensitivity	Human Cysteinyl Leukotriene R1/CysLTR1. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except NMU (45%).
Immunogen	Synthetic 20 amino acid peptide from 3rd extracellular domain of human Cysteinyl Leukotriene R1/CysLTR1.

Product Application Details	
Applications	Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin 6 - 14 ug/ml
Application Notes	FLOW reactivity reported in (PMID:28115217).




Images

Immunohistochemistry-Paraffin: Cysteinyl Leukotriene R1/CysLTR1 Antibody [NLS1317] - Analysis of anti-CYSLTR1 / CYSLT1 antibody with human lymph node, Hodgkins lymphoma.



Immunohistochemistry-Paraffin: Cysteinyl Leukotriene R1/CysLTR1 Antibody [NLS1317] - Analysis of anti-CYSLTR1 / CYSLT1 antibody with human nasal mucosa, allergic rhinitis, eosinophils.



Publications

Geriolda Topi, Shakti Ranjan Satapathy, Souvik Ghatak, Karin Hellman, Fredrik Ek, Roger Olsson, Roy Ehrnström, Marie-Louise Lydrup, Anita Sjölander High Oestrogen receptor alpha expression correlates with adverse prognosis and promotes metastasis in colorectal cancer Cell Communication and Signaling : CCS 2024-03-28 [PMID: 38549115]

Salimi M, Stoger L, Liu W, Go S, et al. Cysteinyl leukotriene E4 activates human group 2 innate lymphoid cells and enhances the effect of prostaglandin D2 and epithelial cytokines. J. Allergy Clin. Immunol. 2017-10-01 [PMID: 28115217] (FLOW, Human)

Woszczek G, Pawliczak R, Qi HY et al. Functional characterization of human cysteinyl leukotriene 1 receptor gene structure. J Immunol 2005-10-15 [PMID: 16210619] (FLOW, Human)

Procedures

Immunohistochemistry Protocol for CYSLTR1 Antibody (NLS1317)

Immunohistochemistry Protocol for CYSLTR1 Antibody (NLS1317):

Immunohistochemistry

1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
2. Make 4 um sections and place on pre-cleaned and charged microscope slides.
3. Heat in a tissue-drying oven for 45 minutes @ 60 degrees Celcius.
4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene for 5 minutes each @ RT.
5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol for 3 minutes each @ RT.
6. Wash the slides in 2 changes of 95% alcohol for 3 minutes each @ RT.
7. Wash the slides in 1 change of 80% alcohol for 3 minutes @ RT.
8. Rinse the slides in gentle running distilled water for 5 minutes @ RT.
9. Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
11. Rinse the slides in 1X TBS-T for 1 minute @ RT.

****Do not allow the tissues to dry at any time during the staining procedure****

12. Begin the immunostaining by applying a universal protein block for 20 minutes @ RT.
13. Drain protein block from the slides and apply the diluted primary antibody for 45 minutes @ RT.
14. Rinse the slide in 1X TBS-T for 1 minute @ RT.
15. Apply a biotinylated anti-rabbit IgG (H+L) secondary for 30 minutes @ RT.
16. Rinse the slide in 1X TBS-T for 1 minute @ RT.
17. Apply an alkaline phosphatase streptavidin for 30 minutes @ RT.
18. Rinse the slide in 1X TBS-T for 1 minute @ RT.
19. Apply an alkaline phosphatase chromagen substrate for 30 minutes @ RT.
20. Rinse the slide in distilled water for 1 minute @ RT.

****This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)****

21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol for 1 minute each @ RT.
22. Wash the slides in 2 changes of 95% alcohol for 1 minute each @ RT.
23. Wash the slides in 3 changes of 100% alcohol for 1 minute each @ RT.
24. Wash the slides in 3 changes of xylene for 1 minute each @ RT.
25. Apply cover slip.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NLS1317

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-85356PEP	Cysteinyl Leukotriene R1/CysLTR1 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS1317

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

