

Product Datasheet

FPRL1/FPR2 Antibody - BSA Free NLS1878

Unit Size: 0.05 ml

Store at -20 degrees C. Avoid freeze/thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 4 Publications: 35

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NLS1878

Updated 10/23/2024 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NLS1878



NLS1878

FPRL1/FPR2 Antibody - BSA Free

Product Information	
Unit Size	0.05 ml
Concentration	1.1 mg/ml
Storage	Store at -20 degrees C. Avoid freeze/thaw cycles.
Clonality	Polyclonal
Preservative	0.01% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS, 30% Glycerol
Target Molecular Weight	38 kDa


Product Description	
Host	Rabbit
Gene ID	2358
Gene Symbol	FPR2
Species	Human, Mouse, Rat, Bacteria
Reactivity Notes	Rat reactivity reported in scientific literature (PMID: 24086560). Bacteria reactivity reported in scientific literature (PMID: 31234710).
Immunogen	This FPRL1/FPR2 antibody is made to a synthetic peptide from the human FPRL1 protein (between residues 300-350) [UniProt P25090]

Product Application Details	
Applications	Western Blot, Electron Microscopy, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:1000, Flow Cytometry, Immunohistochemistry 1:300, Immunocytochemistry/ Immunofluorescence 1:20-1:75, Immunohistochemistry-Paraffin 1:300, Immunohistochemistry-Frozen reported in scientific literature (PMID 24086560), Electron Microscopy
Application Notes	In Western Blot, a band is seen ~38 kDa representing FPRL1. In ICC/IF, plasma membrane staining was observed in Raw264.7 cells. In IHC-P, staining was also observed in the plasma membrane of human kidney cancer tissue. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended. Customers have reported success in IF on FFPE mouse kidney tissue, following microwave antigen retrieval.




Images


Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Immunoblot of FPR2 protein in various parts of the rat brain including olfactory bulb (OB), prefrontal cortex (PFC), primary somatosensory cortex (CTX1), parietal association cortex and secondary auditory cortex (CTX2), striatum (STR), thalamus and hypothalamus (THA), hippocampus (HPC), cerebellum (CB), brainstem (BS), cervical spinal cord [SC(C)], thoracic spinal cord [SC(T)], and lumbar spinal cord [SC(L)]. Blots incubated with antigen-absorbed antibody i. e. peptide competition, show reduced band intensities. Image collected and cropped by Citeab from the following publication (Localisation of Formyl-Peptide Receptor 2 in the Rat Central Nervous System and Its Role in Axonal and Dendritic Outgrowth. *Neurochem Res* (2018)) licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Mouse J774A.1, mouse reticulum cell sarcoma macrophage cell line. Left panel is + Anti-Rabbit FITC; middle panel indicates the cell nuclei stained with Hoechst; the right panel is a merged image. ICC/IF image submitted by a verified customer review.




Immunohistochemistry-Paraffin: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - SPM receptors differentially expressed on endothelium and infiltrating leukocytes AE" ALX/FPR2 and ChemR23. Acute inflammation triggered by ventral aspect of forearm of healthy volunteers by intradermal injection of 1.5 A-107 UV-killed E.coli (UVkEc) suspended in 100 μ l of saline. 4hrs after injection a 3-mm skin punch biopsy taken from inflamed site under local anesthesia. Naive skin treated as baseline. IHC-P on skin sections for receptor identification. Low mag(A-5) and high-mag (A-40) images at baseline and the 4 hr time point shown for ALX/FPR2. Red arrows highlight endothelium. Black arrow highlights infiltrating leukocytes. Image collected and cropped by Citeab from the (Pro-resolving mediators promote resolution in human skin model of UV-killed Escherichia coli-driven acute inflammation *JCI Insight* (2018)) licensed under, CC-BY license.



Flow Cytometry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Flow Cytometry: [Alexa Fluor® 700] [NLS1878AF700] - Mouse splenocytes. Cells were pre-gated with live/dead and FSC-A/W to exclude dead cells and cell doublets. Image from verified customer review. Image using the Alexa Fluor 700 form of this antibody.




Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Murine neutrophils were extracted from the bone marrow of C57BL/6 mice and exposed to normal glucose (90 mg/dl) or high glucose (300 mg/dl) and the expression of FPR2 was assessed by western blotting after 1, 2, or 3 hr exposure to glucose. Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/35112667/>) licensed under a CC-BY license.




Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Analysis in HL-60 cell lysate.




Immunocytochemistry/Immunofluorescence: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Antibody was tested in Raw264.7 cells with DyLight 488 (green). Nuclei were counterstained with DAPI (blue).




Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Analysis in human kidney cancer using DAB with hematoxylin counterstain.




Electron Microscopy: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c=50A nm, e=100A nm. Image collected and cropped by Citeab from the following publication (Localisation of Formyl-Peptide Receptor 2 in the Rat Central Nervous System and Its Role in Axonal and Dendritic Outgrowth. *Neurochem Res* (2018)) licensed under a CC-BY license.




Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] -
 Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Electron micrographs of FPR2 immunostained sections from the prefrontal cortex. a, b Immunostaining is mostly present in axon pre-terminals (PrT) that did not form synapses with postsynaptic structures. c Occasional axon terminals (AT) are observed to form asymmetrical, putatively glutamatergic synapses (S) with unlabelled dendrites (DE). d Occasional labelled dendrites (DE) are also found, that formed asymmetrical synapses (S) with unlabelled axon terminals (AT). Scale: a, b, c = 50 nm, e = 100 nm Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29948727>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Lipoxin A4, along with its precursors & ligand, were altered by hAECs. (A): Expression of the lipoxin A4 precursor genes ALOX ζ 5, ζ 12, & ζ 15 was increased at days 5 & 7 compared with saline controls (ALOX ζ 5: 0.393 ± 0.114 vs. 3.777 ± 1.98 ; ALOX ζ 12: 0.484 ± 0.248 vs. 1.520 ± 0.203 ; ALOX ζ 15: 0.020 ± 0.007 vs. 1.998 ± 0.983). \square , $p < .05$. (B): Lipoxin A4 protein levels were elevated in lung lysates at day 7 in animals treated with hAECs compared with controls (0.103 ± 0.021 ng/ml vs. 0.249 ± 0.072 ng/ml, respectively). \square , $p < .05$. (C): At day 7, bleomycin challenge resulted in positively stained F4/80/FPR2 cells, which was elevated in mice treated with hAECs compared with control mice ($5.720\% \pm 0.587\%$ vs. $8.795\% \pm 0.687\%$, respectively). \square , $p < .05$. (D): Representative images of F4/80 \square & FPR2 \square positive \square stained lung sections from hAEC \square treated animals. Magnification: $\times 200$. Scale bar = $100 \mu\text{m}$. Abbreviations: DAPI, 4',6-diamidino-2-phenylindole; FPR2, N-formyl peptide receptor 2; hAEC, human amnion epithelial cell. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/28371562>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: FPRL1/FPR2 Antibody - BSA Free [NLS1878] - Higher expression of Fpr2 in the livers of female mice is related with hepatocyte protection. a qRT-PCR analysis for Fpr2 expression in primary hepatocytes (pHEPs) from WT (male $n = 2$, female $n = 2$) & KO mice (male $n = 2$, female $n = 2$). Total eight mice were employed in each hepatocyte isolation, & the experiments were replicated at least three times & the mean \pm S.E.M. results are graphed ($*p < 0.05$, $**p < 0.005$ vs WT male-pHEPs). b Western blot analysis & c double immunofluorescent images of Fpr2 (red) with albumin (green) in these cells. Gapdh was used as internal control. DAPI (blue) was used as nuclear counterstaining. Data shown represent one of three experiments with similar results (Scale bar, $20 \mu\text{m}$). d qRT-PCR analysis for Fpr2 & glucose-6-phosphatase (G6pc) in, e cell viability of, & f western blot analysis of cleaved Caspase-3 & pro Caspase-3 in WT & KO female mice-isolated pHEPs treated with vehicle (Veh) or $250 \mu\text{M}$ of palmitate (PA). The data shown represent one of three experiments with similar results. The mean \pm S.E.M. results obtained from three repetitive experiments are graphed ($*p < 0.05$, $**p < 0.005$ vs WT-Veh). Gray circles represent individual data points. See Supplementary Data for statistical details. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/35102146>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Zhu M, Wang Y, Zhu L et al. Crosstalk between RPE cells and choroidal endothelial cells via the ANXA1/FPR2/SHP2/NLRP3 inflammasome/pyroptosis axis promotes choroidal neovascularization Inflammation 2021-10-01 [PMID: 34595678]

Federica Frigerio, Giulia Pasqualini, Ilaria Craparotta, Sergio Marchini, Erwin A van Vliet, Patrick Foerch, Catherine Vandenas, Karin Leclercq, Eleonora Aronica, Luca Porcu, Kimberly Pistorius, Romain A Colas, Trond V Hansen, Mauro Perretti, Rafal M Kaminski, Jesmond Dalli, Annamaria Vezzani n-3 Docosapentaenoic acid-derived protectin D1 promotes resolution of neuroinflammation and arrests epileptogenesis Brain 2018-11-01 [PMID: 30307467]

R Roy, J Zayas, SK Singh, K Delgado, SJ Wood, MF Mohamed, DM Frausto, YA Albalawi, TP Price, R Estupinian, EF Giurini, TM Kuzel, A Zloza, J Reiser, SH Shafikhani Overriding impaired FPR chemotaxis signaling in diabetic neutrophil stimulates infection control in murine diabetic wound Elife, 2022-02-03;11(0):. 2022-02-03 [PMID: 35112667]

YE Kim, SY Ahn, DK Sung, YS Chang, WS Park Mesenchymal Stem Cells and Formyl Peptide Receptor 2 Activity in Hyperoxia-Induced Lung Injury in Newborn Mice International Journal of Molecular Sciences, 2022-09-13;23(18):. 2022-09-13 [PMID: 36142517]

C Lee, J Kim, J Han, D Oh, M Kim, H Jeong, TJ Kim, SW Kim, JN Kim, YS Seo, A Suzuki, JH Kim, Y Jung Formyl peptide receptor 2 determines sex-specific differences in the progression of nonalcoholic fatty liver disease and steatohepatitis Nature Communications, 2022-01-31;13(1):578. 2022-01-31 [PMID: 35102146]

Studley WR, Lamanna E, Martin KA et al. The small-molecule formyl peptide receptor biased agonist, compound 17b, is a vasodilator and anti-inflammatory in mouse precision-cut lung slices British journal of pharmacology 2023-09-01 [PMID: 37658546] (IHC-P, Mouse)

Li L, Cheng SQ, Sun YQ et al. Resolvin D1 reprograms energy metabolism to promote microglia to phagocytize neutrophils after ischemic stroke Cell reports 2023-06-06 [PMID: 37285269] (IHC, WB, Mouse)

Liu M, He H, Fan F et al. Maresin-1 protects against pulmonary arterial hypertension by improving mitochondrial homeostasis through ALXR/HSP90 α axis Journal of molecular and cellular cardiology 2023-05-25 [PMID: 37244057] (WB, Rat)

Liu L, Kim S, Buckley MT et al. Exercise reprograms the inflammatory landscape of multiple stem cell compartments during mammalian aging Cell stem cell 2023-04-13 [PMID: 37080206] (IHC, Mouse)

Tang D, Fu G, Li W et al. Maresin 1 Protects the Liver Against Ischemia/Reperfusion Injury via the ALXR/Akt Signaling Pathway Mol Med 2021-02-26 [PMID: 33632134]

Kim SY, Kim JM, Lee SR et al. Efferocytosis and enhanced FPR2 expression following apoptotic cell instillation attenuate radiation-induced lung inflammation and fibrosis Biochemical and biophysical research communications 2022-02-21 [PMID: 35228119] (WB)

Jun JH, Park S, Kim JY et al. Combination Therapy of Placenta-Derived Mesenchymal Stem Cells with WKYMVm Promotes Hepatic Function in a Rat Model with Hepatic Disease via Vascular Remodeling Cells 2022-01-11 [PMID: 35053347] (ICC/IF)

More publications at <http://www.novusbio.com/NLS1878>

Procedures

Immunohistochemistry-Paraffin protocol for FPRL1 Antibody (NLS1878)

Immunohistochemistry-Paraffin Embedded Sections

Antigen Unmasking:

Bring slides to a boil in 10 mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench-top for 30 minutes.

Staining:

1. Wash sections in deionized water three times for 5 minutes each.
2. Wash sections in wash buffer for 5 minutes.
3. Block each section with 100-400 ul blocking solution for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul diluted primary antibody. Incubate overnight at 4 C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated diluted secondary antibody. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in deionized water.
12. Counterstain sections in hematoxylin.
13. Wash sections in deionized water two times for 5 minutes each.
14. Dehydrate sections.
15. Mount coverslips.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.

Western Blot protocol for FPRL1 Antibody (NLS1878)

Western Blot Protocol

1. Perform SDS-PAGE on samples to be analyzed, loading 40 ug of total protein per lane.
2. Transfer proteins to membrane according to the instructions provided by the manufacturer of the membrane and transfer apparatus.
3. Stain according to standard Ponceau S procedure (or similar product) to assess transfer success, and mark molecular weight standards where appropriate.
4. Rinse the blot.
5. Block the membrane using standard blocking buffer for at least 1 hour.
6. Wash the membrane in wash buffer three times for 10 minutes each.
7. Dilute primary antibody in blocking buffer and incubate 1 hour at room temperature.
8. Wash the membrane in wash buffer three times for 10 minutes each.
9. Apply the diluted HRP conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer three times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions.

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



Immunocytochemistry/Immunofluorescence Protocol for FPRL1 Antibody (NLS1878)

Immunocytochemistry Protocol

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and wash the cells briefly in PBS. Add 10% formalin to the dish and fix at room temperature for 10 minutes.
2. Remove the formalin and wash the cells in PBS.
3. Permeabilize the cells with 0.1% Triton X100 or other suitable detergent for 10 min.
4. Remove the permeabilization buffer and wash three times for 10 minutes each in PBS. Be sure to not let the specimen dry out.
5. To block nonspecific antibody binding, incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
6. Add primary antibody at appropriate dilution and incubate overnight at 4C.
7. Remove primary antibody and replace with PBS. Wash three times for 10 minutes each.
8. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
9. Remove secondary antibody and replace with PBS. Wash three times for 10 minutes each.
10. Counterstain DNA with DAPI if required.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NLS1878

NLS1878PEP	FPRL1/FPR2 Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS1878

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

