Product Datasheet

P2Y12/P2RY12 Antibody - BSA Free NLS3742

Unit Size: 0.05 ml

Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NLS3742

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NLS3742



NLS3742


P2Y12/P2RY12 Antibody - BSA Free

| PZY12/PZRY1Z ANIIDOGY - BSA Free | |
|----------------------------------|---|
| Product Information | |
| Unit Size | 0.05 ml |
| Concentration | 1.0 mg/ml |
| Storage | Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles. |
| Clonality | Polyclonal |
| Preservative | 0.1% Sodium Azide |
| Isotype | IgG |
| Purity | Immunogen affinity purified |
| Buffer | PBS |
| Product Description | |
| Description | Product can be stored undiluted at 4C for up to 1 month. |
| Host | Rabbit |
| Gene ID | 64805 |
| Gene Symbol | P2RY12 |
| Species | Human |
| Reactivity Notes | Predicted cross-reactivity based on sequence identity: Gorilla (100%), Gibbon (88%), Marmoset (88%), Porcine (82%). |
| Specificity/Sensitivity | Human P2Y12/P2RY12. BLAST analysis of the peptide immunogen showed no homology with other human proteins. |
| Immunogen | Synthetic 17 amino acid peptide from N-terminal extracellular domain of human P2Y12/P2RY12. |
| Product Application Details | |
| Applications | Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin |
| Recommended Dilutions | Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, |

Immunohistochemistry-Paraffin 32 ug/ml


Images

Immunocytochemistry/Immunofluorescence: P2Y12/P2RY12 Antibody [NLS3742] - Analysis of anti-P2RY12 / P2Y12 antibody with HEK293 human embryonic kidney cells







Immunohistochemistry-Paraffin: P2Y12/P2RY12 Antibody [NLS3742] - Ovary, Carcinoma




Immunohistochemistry-Paraffin: P2Y12/P2RY12 Antibody [NLS3742] - Analysis of anti-P2RY12 / P2Y12 antibody with human brain, glioblastoma.




Immunohistochemistry-Paraffin: P2Y12/P2RY12 Antibody [NLS3742] - Analysis of anti-P2RY12 / P2Y12 antibody with human blood, platelets.



Immunohistochemistry-Paraffin: P2Y12/P2RY12 Antibody [NLS3742] - Breast, Carcinoma



Immunohistochemistry-Paraffin: P2Y12/P2RY12 Antibody [NLS3742] - Lung, Small Cell Carcinoma



Procedures

Immunohistochemistry Protocol for P2Y12 Antibody (NLS3742)

Immunohistochemistry Protocol for P2Y12 Antibody (NLS3742): Immunohistochemistry

- 1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
- 2. Make 4-um sections and place on pre-cleaned and charged microscope slides.
- 3. Heat in a tissue-drying oven for 45 minutes at 60 degrees Celcius.
- 4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene approximately 5 minutes each @ RT.
- 5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol approximately 3 minutes each @ RT.
- 6. Wash the slides in 2 changes of 95% alcohol approximately 3 minutes each @ RT.
- 7. Wash the slides in 1 change of 80% alcohol approximately 3 minutes @ RT.
- 8. Rinse the slides in gentle running distilled water approximately 5 minutes @ RT.
- 9. Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
- 10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
- 11. Rinse the slides in 1X TBS-T for 1 minute @ RT.
- **Do not allow the tissues to dry at any time during the staining procedure**
- 12. Begin the immunostaining by applying a universal protein block approximately 20 minutes @ RT.
- 13. Drain protein block from the slides and apply the diluted primary antibody approximately 45 minutes @ RT.
- 14. Rinse the slide in 1X TBS-T approximately 1 minute @ RT.
- 15. Apply a biotinylated anti-rabbit IgG (H+L) secondary approximately 30 minutes @ RT.
- 16. Rinse the slide in 1X TBS-T approximately 1 minute at RT.
- 17. Apply an alkaline phosphatase steptavidin approximately 30 minutes at RT.
- 18. Rinse the slide in 1X TBS-T approximately 1 minute at RT.
- 19. Apply an alkaline phosphatase chromagen substrate approximately 30 minutes at RT.
- 20. Rinse the slide in distilled water approximately 1 minute @ RT.
- **This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)**
- 21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol approximately 1 minute each @ RT.
- 22. Wash the slides in 2 changes of 95% alcohol approximately 1 minute each @ RT.
- 23. Wash the slides in 3 changes of 100% alcohol approximately 1 minute each @ RT.
- 24. Wash the slides in 3 changes of xyleneapproximately 1 minute each @ RT.
- 25. Apply cover slip.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NLS3742

NBL1-14038 P2Y12/P2RY12 Overexpression Lysate

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS3742

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

