

Product Datasheet

Histamine H4R Antibody - BSA Free NLS3775

Unit Size: 0.05 ml

Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NLS3775

Updated 4/1/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NLS3775



NLS3775

Histamine H4R Antibody - BSA Free

Product Information

Unit Size	0.05 ml
Concentration	1.0 mg/ml
Storage	Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS

Product Description


Description	Product can be stored undiluted at 4C for up to 1 month.
Host	Rabbit
Gene ID	59340
Gene Symbol	HRH4
Species	Human, Monkey
Reactivity Notes	Predicted cross-reactivity based on sequence identity: Marmoset (100%), Gibbon (89%), Mouse (89%), Canine (89%), Rat (83%), Opossum (83%).
Specificity/Sensitivity	Human Histamine H4R. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Immunogen	Synthetic 18 amino acid peptide from 1st cytoplasmic domain of human Histamine H4R.

Product Application Details


Applications	Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 20 ug/ml
Application Notes	.

Images


Immunohistochemistry-Paraffin: Histamine H4R Antibody [NLS3775] - Lung, Non Small-Cell Carcinoma




Immunohistochemistry-Paraffin: Histamine H4 R Antibody [NLS3775] - IHC analysis of Lymph node using this antibody at 12 ug/ ml.




Immunohistochemistry-Paraffin: Histamine H4 R Antibody [NLS3775] - Anti-HRH4 / Histamine H4 Receptor antibody IHC of human Ovary, Carcinoma and Inflammatory Cells. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.




Immunohistochemistry-Paraffin: Histamine H4R Antibody [NLS3775] - Ovary, carcinoma




Immunohistochemistry-Paraffin: Histamine H4R Antibody [NLS3775] - Pancreas, carcinoma



Immunohistochemistry-Paraffin: Histamine H4R Antibody [NLS3775] - Colon, Carcinoma



Immunohistochemistry-Paraffin: Histamine H4R Antibody - BSA Free [NLS3775] - Immunohistochemistry of formalin-fixed, paraffin-embedded Human Testis, Mast Cells tissue



Publications

S Moon, O Stasikowska-Kanicka, M Wągrowaska-Danilewicz, M Hawro, M Metz, M Maurer, T Hawro Clinically uninvolved but not healthy-The skin of patients with atopic dermatitis is primed for itch and inflammation. Journal of the European Academy of Dermatology and Venereology : JEADV 2023-12-08 [PMID: 38063249]

Wang Y Analysis of mast cells and mast cell-mediator-related histological features in cholinergic urticaria Thesis (IF/IHC, Human)

Procedures

Immunohistochemistry Protocol for Histamine H4 Receptor Antibody (NLS3775)

Immunohistochemistry Protocol for Histamine H4 Receptor Antibody (NLS3775):

Immunohistochemistry

1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
2. Make 4 um sections and place on pre-cleaned and charged microscope slides.
3. Heat in a tissue-drying oven for 45 minutes @ 60 degrees Celcius.
4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene for 5 minutes each @ RT.
5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol for 3 minutes each @ RT.
6. Wash the slides in 2 changes of 95% alcohol for 3 minutes each @ RT.
7. Wash the slides in 1 change of 80% alcohol for 3 minutes @ RT.
8. Rinse the slides in gentle running distilled water for 5 minutes @ RT.
9. Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
11. Rinse the slides in 1X TBS-T for 1 minute @ RT.

****Do not allow the tissues to dry at any time during the staining procedure****

12. Begin the immunostaining by applying a universal protein block for 20 minutes @ RT.
13. Drain protein block from the slides and apply the diluted primary antibody for 45 minutes @ RT.
14. Rinse the slide in 1X TBS-T for 1 minute @ RT.
15. Apply a biotinylated anti-rabbit IgG (H+L) secondary for 30 minutes @ RT.
16. Rinse the slide in 1X TBS-T for 1 minute @ RT.
17. Apply an alkaline phosphatase streptavidin for 30 minutes @ RT.
18. Rinse the slide in 1X TBS-T for 1 minute @ RT.
19. Apply an alkaline phosphatase chromagen substrate for 30 minutes @ RT.
20. Rinse the slide in distilled water for 1 minute @ RT.

****This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)****

21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol for 1 minute each @ RT.
22. Wash the slides in 2 changes of 95% alcohol for 1 minute each @ RT.
23. Wash the slides in 3 changes of 100% alcohol for 1 minute each @ RT.
24. Wash the slides in 3 changes of xylene for 1 minute each @ RT.
25. Apply cover slip.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NLS3775

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
210-TA-005	TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS3775

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

