Product Datasheet

alpha-1B Adrenergic R/ADRA1B Antibody - BSA Free NLS697

Unit Size: 0.05 ml

Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NLS697

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NLS697



NLS697

alpha-1B Adrenergic R/ADRA1B Antibody - BSA Free

Product Information	
0.05 ml	
1.0 mg/ml	
Keep as concentrated solution. Aliquot and store at -20C or below. Avoid multiple freeze-thaw cycles.	
Polyclonal	
0.1% Sodium Azide	
IgG	
Immunogen affinity purified	
PBS	
Product Description	
Product can be stored undiluted at 4C for up to 1 month.	
Rabbit	
147	
ADRA1B	
Human, Mouse, Rat, Bovine, Canine, Hamster, Monkey, Primate	
Predicted cross-reactivity based on sequence identity: Gorilla (100%), Marmoset (100%), Bat (85%), Opossum (80%), Platypus (80%).	

Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin 11 ug/ml

Human ADRA1B. BLAST analysis of the peptide immunogen showed no

Synthetic 20 amino acid peptide from C-terminus of human alpha-1B Adrenergic

homology with other human proteins, except PLEKHG5 (45%).


Images

Immunogen

Specificity/Sensitivity

Immunohistochemistry-Paraffin: alpha-1B Adrenergic R/ADRA1B Antibody [NLS697] - Analysis of anti-ADRA1B antibody with human adrenal, medulla.

R/ADRA1B.



Publications

Karabacakb Or, Sener Nc, Yilmazer D. Alpha Adrenergic Receptors In Renal Pelvis And Calyces: Can Rat Models Be Used? Int Braz J Urol 2014-12-16 [PMID: 25498280] (IHC-P, Rat)

Williams LM, Qi Z, Batai K et al. A locus on chromosome 5 shows African ancestry-limited association with alloimmunization in sickle cell disease Blood Adv 2018-12-26 [PMID: 30578281] (IHC-P, Human)

Karabacak OR, Yilmazer D, Ozturk U et al. The presence and distribution of alpha adrenergic receptors in human renal pelvis and calyces. Urolithiasis. 2013-10-01 [PMID: 23877383]

Details:

ADRA 1A (IMG-71129), ADRA1B (IMG-71127) and ADRA 1D (IMG-71130).



Procedures

Immunohistochemistry Protocol for Alpha 1b Adrenergic Receptor Antibody (NLS697)

Immunohistochemistry Protocol for Alpha 1b Adrenergic Receptor Antibody (NLS697): Immunohistochemistry

- 1. Prepare tissue with formalin fixation and by embedding it in paraffin wax.
- 2. Make 4 um sections and place on pre-cleaned and charged microscope slides.
- 3. Heat in a tissue-drying oven for 45 minutes @ 60 degrees Celcius.
- 4. Deparaffinize the tissues by wash drying the slides in 3 changes of xylene for 5 minutes each @ RT.
- 5. Rehydrate the tissues by washing the slides in 3 changes of 100% alcohol for 3 minutes each @ RT.
- 6. Wash the slides in 2 changes of 95% alcohol for 3 minutes each @ RT.
- 7. Wash the slides in 1 change of 80% alcohol for 3 minutes @ RT.
- 8. Rinse the slides in gentle running distilled water for 5 minutes @ RT.
- 9. Perform antigen retrieval by steaming the slides in 0.01M sodium citrate buffer (pH 6.0) @ 99-100 degrees Celcius for 20 minutes.
- 10. Remove the slides from the heat and let stand in buffer @ RT for 20 minutes.
- 11. Rinse the slides in 1X TBS-T for 1 minute @ RT.
- **Do not allow the tissues to dry at any time during the staining procedure**
- 12. Begin the immunostaining by applying a universal protein block for 20 minutes @ RT.
- 13. Drain protein block from the slides and apply the diluted primary antibody for 45 minutes @ RT.
- 14. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 15. Apply a biotinylated anti-rabbit IgG (H+L) secondary for 30 minutes @ RT.
- 16. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 17. Apply an alkaline phosphatase steptavidin for 30 minutes @ RT.
- 18. Rinse the slide in 1X TBS-T for 1 minute @ RT.
- 19. Apply an alkaline phosphatase chromagen substrate for 30 minutes @ RT.
- 20. Rinse the slide in distilled water for 1 minute @ RT.
- **This method should only be used if the chromagen substrate is alcohol insoluble (ie: Vector Red, DAB)**
- 21. Dehydrate the tissue by washing the slides in 2 changes of 80% alcohol for 1 minute each @ RT.
- 22. Wash the slides in 2 changes of 95% alcohol for 1 minute each @ RT.
- 23. Wash the slides in 3 changes of 100% alcohol for 1 minute each @ RT.
- 24. Wash the slides in 3 changes of xylene for 1 minute each @ RT.
- 25. Apply cover slip.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NLS697

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-57288PEP alpha-1B Adrenergic R/ADRA1B Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NLS697

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

